



Medicinskt
programarbete

Referensförteckning

Regionalt vårdprogram

Depression
och bipolär sjukdom

Stockholms läns landsting

2007





Referensförteckning

Regionalt vårdprogram

Depression
och bipolär sjukdom

Rapporten är framtagen av
Christina Spjut

ISBN 91-85211-45-1
(Reviderad utgåva 2003, ISBN 91-85209-14-7)

 **Stockholms läns landsting**
Forum för Kunskap och gemensam Utveckling

Innehåll

Epidemiologi och könets betydelse för affektiv sjukdom.....	5
Hälsoekonomi	6
Prevention	6
Begrepp och definitioner	7
Depression och suicid	8
Diagnostik och skattning	11
Läkemedelsbehandling av depression	12
Behandling av bipolär sjukdom	12
Psykologisk behandling vid depression och bipolär sjukdom	12
Fysikaliska behandlingsmetoder.....	15
Omvårdnad	15
Den terapeutiska alliansen	17
Affektiv sjukdom under graviditet och postpartum.....	17
Bipolär sjukdom hos barn och unga	22
Affektiv sjukdom hos personer över 65 år.....	22

Epidemiologi och könets betydelse för affektiv sjukdom

1. Kessler, R.C., et al., Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States. Results from the National Comorbidity Survey. *Arch Gen Psychiatry*, 1994. 51(1): p. 8-19.
2. Keller, 2001
3. Yorbik, O., et al., Clinical characteristics of depressive symptoms in children and adolescents with major depressive disorder. *J Clin Psychiatry*, 2004. 65(12): p. 1654-9; quiz 1760-1.
4. Christensen, E.M., et al., Life events and onset of a new phase in bipolar affective disorder. *Bipolar Disord*, 2003. 5(5): p. 356-61.
5. Baldassano, C. F., L. B. Marangell, et al. (2005). "Gender differences in bipolar disorder: retrospective data from the first 500 STEP-BD participants." *Bipolar Disord* 7(5): 465-70.
6. Kendler, K.S., J. Myers, and C.A. Prescott, Sex differences in the relationship between social support and risk for major depression: a longitudinal study of opposite-sex twin pairs. *Am J Psychiatry*, 2005. 162(2): p. 250-6.
7. Cahill, L., His brain, her brain. *Sci Am*, 2005. 292(5): p. 40-7.
8. Raskin, A., Age-sex differences in response to antidepressant drugs. *J Nerv Ment Dis*, 1974. 159(2): p. 120-30.
9. Davidson, J. and S. Pelton, Forms of atypical depression and their response to antidepressant drugs. *Psychiatry Res*, 1986. 17(2): p. 87-95.
10. Quitkin, F.M., et al., Are there differences between women's and men's antidepressant responses? *Am J Psychiatry*, 2002. 159(11): p. 1848-54.
11. Kornstein, S.G., et al., Gender differences in treatment response to sertraline versus imipramine in chronic depression. *Am J Psychiatry*, 2000. 157(9): p. 1445-52.

12. Baca, E., M. Garcia-Garcia, and A. Porrás-Chavarino, Gender differences in treatment response to sertraline versus imipramine in patients with nonmelancholic depressive disorders. *Prog Neuropsychopharmacol Biol Psychiatry*, 2004. 28(1): p. 57-65.
13. Grigoriadis, S., S.H. Kennedy, and R.M. Bagby, A comparison of antidepressant response in younger and older women. *J Clin Psychopharmacol*, 2003. 23(4): p. 405-7. 24. Thase, M.E., et al., Gender differences in response to treatments of depression, in *Gender and its effect on psychopathology*, F.E. Washington, Editor. 2000, American Psychiatric Association Press: Washington DC. p. 103-129.

Hälsoekonomi

1. Sobocki, 2006

Prevention

1. Folkhälsoinstitutet. Att överbrygga hälsogapen i Sverige genom hälsofrämjande och sjukdomsförebyggande arbete. Folkhälsoinstitutet 1996:90. Stockholm.
2. Jarman B, Hirsch S, White P, Driscoll R. Predicting psychiatric admission rates. *B M J* 1992;304:1146-1151.
3. Alonso J, Codony M, Kovess V, Angermeyer MC, Katz SJ, Haro JM et al. Population level of unmet need for mental care in Europe. *Br J Psychiatry* 2007;190:299-306
4. Dahlberg K, Forsell Y, Damström-Thakker K, Runeson B. Mental health problems healthcare contacts in an urban and a rural area. Comparisons of two Swedish counties. *Nord J Psychiatry* 2007;61:40-46.
5. Forsell Y. The pathway to meeting need for mental health services in Sweden. *Psychiatric Services* 2006;57:114-119.
6. Psykisk hälsa. Rapport från arbetsgruppen för psykisk hälsa till Nationella folkhälsokommittén. Underlagsrapport nr 10. Stockholm 1999.
7. Maslach C. Workshop on Burnout and Depression. Nobel Forum, Karolinska Institutet, Stockholm 2001.

8. Maslach C, Shaufeli WB, Leiter MP. Job burnout. *Annu Rev Psychol* 2001:
9. Katon W, Sullivan M. Depression and Chronic Medical Illness. *Journal of Clinical Psychiatry* 1990;5:3-11.
10. Ekman M. Ensamma mammor – en grupp utsatt för särskilda hälsorisker. Ur: Att överbygga hälsogapen i Sverige genom hälsofrämjande och sjukdomsförebyggande arbete. *Folkhälsoinstitutet* 1996;90:221-233.
11. Schoevers R.A. et al Prevention of Late-Life Depression in Primary Care: Do We Know Where to Begin? *Am J Psychiatry* 2006; 163:1611-162
12. Rutz W, von Knorring, Wålinder J. Frequency of suicide on Gotland after systematic postgraduate education of general practitioners. *Acta Psychiatr Scand* 1989;80:151-154.
13. Rutz W, von Knorring L, Wålinder J, Wistedt B. Effect of an educational program for general practitioners on Gotland on the pattern of prescription of psychotropic drugs. *Acta Psychiatr Scand* 1990;82:399-403.

Begrepp och definitioner

1. Angst, 1992
2. Angst, 1986
3. Akiskal, 1996
4. Lopez-Ibor, 1994
5. Burch, 1994
6. Post, 1989

Depression och suicid

1. Guze SB, Robins E. Suicide and primary affective disorders. *Brit J Psychiatry* 1970;117:437-438.
2. Bostwick JM, Pankratz VS. Affective disorders and suicide risk: A re-examination. *Am J Psychiatry* 2000;157:1925-1932
3. Ösby U, Brandt L, Correia N, Ekblom A, Sparen P. Excess mortality in bipolar and unipolar disorder in Sweden. *Arch Gen Psychiatry* 2001 Sep;58(9):844-50
4. Harris EC, Barraclough B. Suicide as an outcome for mental disorders. A meta-analysis. *Brit J Psychiatry* 1997;170:205-228
5. Angst J, Angst F, Gerber-Werder R, Gamma A. Suicide in 406 mood-disorder patients with and without long-term medication: a 40 to 44 years' follow-up. *Arch Suicide Res* 2005;9(3):279-300
6. Barraclough B, Bunch J, Nelson B. A hundred cases of suicide: clinical aspects. *Br J Psychiatry* 1974;125:355-374
7. Barraclough BM, Pallis DJ. Depression followed by suicide: a comparison of depressed suicides with living depressives. *Psychol Med* 1975 Feb;5(1):55-61
8. Cheng A. Mental illness and suicide. A case-control study in East Taiwan. *Arch Gen Psychiatry* 1995;52:594-603
9. Rifai AH, George CJ, Stack JA, Mann JJ, Reynolds CF Hopelessness in suicide attempters after acute treatment of major depression in late life. *Am J Psychiatry* 1994;151(11):1687-90
10. Jick SS, Dean AD, Jick H. Antidepressants and suicide. *Brit Med J* 1995;28;310:215-8
11. Fawcett J, Scheftner WA, Fogg L, et al. Time-related predictors of suicide in major affective disorder. *Am J Psychiatry* 1990;147:1189-1194.
12. Busch KA, Fawcett J, Jacobs DG. Clinical correlates of inpatient suicide. *J Clin Psychiatry* 2003;64:14-19.
13. Agargun MY, Kara H, Solmaz M. Sleep disturbances and suicidal behaviour in patients with major depression. *J Clin Psychiatry* 1997;58:249-251.

14. Agargun MY, Kara H, Solmaz M. Subjective sleep quality and suicidality in patients with major depression. *J Psychiatric Res* 1997;3:377-381.
15. Jamison K. Suicide in bipolar disorder. Föredrag vid APA, New Orleans 2001.
16. Stone MH. Long-term outcome in personality disorders. *Brit J Psychiatry* 1993;162:299-313.
17. Runeson B, Beskow J, Waern M. The suicidal process in suicides among young people. *Acta Psychiatr Scand* 1996;93:35-42.
18. Heilä H, Isometsä ET, Henriksson MM, Heikkinen ME, Marttunen MJ, Lönnqvist JK. Suicide and schizophrenia: A nationwide psychological autopsy study on age- and sex-specific clinical characteristics of 92 suicide victims with schizophrenia. *Am J Psychiatry* 1997;154:1235-1242.
19. Heilä H, Isometsä ET, Henriksson MM, Heikkinen ME, Marttunen MJ, Lönnqvist JK: Antecedents of suicide in people with schizophrenia. *Brit J Psychiatry* 1998; 173: 330-3.
20. Berglund M, Öjehagen A. The influence of alcohol drinking and alcohol use disorders on psychiatric disorders and suicidal behavior. *Alcoholism, Clinical & Experimental Research* 1998;22(7 Suppl S):333S-345S.
21. Runeson B, Beskow J. Borderline personality disorder in young Swedish suicides. *J Nerv Ment Dis* 1991;179(3):153-6.
22. Waern M, Rubenowitz E, Runeson B, Skoog I, Wilhelmsson K, Allebeck P. Burden of illness and suicide in elderly people: case – central study. *Brit Med J* 2002;324:1355-1357.
23. Nordström P, Åsberg M, Åberg-Wistedt A, Nordin C. Attempted suicide predicts suicide risk in mood disorders. *Acta Psychiatr Scand.* 1995;92(5):345-50
24. Rynestad T. Mortality after deliberate self-poisoning. A prospective follow-up study of 587 persons observed for 5279 person years: risk factors and causes of death. *Soc Psych Psychiatr Epidemiol.* 1997 Nov;32:443-50.
25. Tejedor MC, Diaz A, Castillon JJ, Pericay JM. Attempted suicide: repetition and survival-findings of a follow-up study. *Acta Psychiatr Scand.* 1999;100:205-11

26. Appleby L, Cooper J, Amos T, Faragher B. Psychological autopsy study of suicides by people aged under 35. *Br J Psychiatry*. 1999 Aug;175:168-74
27. Jenkins GR, Hale R, Papanastassiou M, Crawford MJ, Tyrer P. Suicide rate 22 years after parasuicide: cohort study. *Brit Med J* 2002; 325: 1155
28. Suominen K, Isometsä E, Suokas J, et al. Completed suicide after a suicide attempt: a 37-year follow-up study. *Am J Psychiatry* 2004;161:563-564
29. Henriksson MM, Isometsä ET, Hietanen PS, Aro HM, Lönnqvist JK. Mental disorders in cancer suicides. *J Affect Dis* 1995;36:11-20.
30. Waern M, Runeson B, Allebeck P, Beskow J, Rubenowitz E, Skoog I, Wilhelmsson K. Mental disorder in elderly suicides: a case-control study. *Am J Psychiatry* 2002;159(3):450-455.
31. Tiihonen J, Lönnqvist J, Wahlbeck K, Klaukka T, Tanskanen A, Haukka J. Antidepressants and the risk of suicide, attempted suicide, and overall mortality in a nationwide cohort. *Arch Gen Psychiatry* 2006;63:1358-1367
32. Goodwin FK, Fireman B, Simon GE, Hunkeler EM, Lee J, Revicki D. Suicide risk in bipolar disorder during treatment with lithium and divalproex. *JAMA* 2003;290:1467-1473.
33. Fergusson D, Doucette S, Glass KC, Shapoiro S, Healy D, Hebert P, Hutton B. Association between suicide attempts and selective serotonin reuptake inhibitors: systematic review of randomised controlled trials. *Brit Med J*. 2005 Feb 19;330(7488):396
34. Gunnell D, Saperia J, Ashby D. Selective serotonin reuptake inhibitors (SSRIs) and suicide in adults: meta-analysis of drug company data from placebo controlled, randomised controlled trials submitted to the MHRA's safety review. *Brit Med J*. 2005;19:330(7488):385
35. Khan A, Khan S, Kolts R, Brown WA. Suicide rates in clinical trials of SSRIs, other antidepressants, and placebo: analysis of FDA reports. *Am J Psychiatry*. 2003 Apr;160(4):790-2.

Diagnostik och skattning

1. APA, Diagnostiska kriterier enligt DSM-IV (Quick reference to the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV). 1995, Danderyd: Pilgrim Press.
2. First, M., et al., Structured Clinical Interview for DSM-IV, Axis I Disorders (SCID-I). Clinical version. Administration booklet (Svensk version: Strukturerad klinisk intervju för DSM-IV axel I. Pilgrim Press, Danderyd). 1998, New York: Biometrics Research Department, New York State Psychiatric Institute.
3. Sheehan, D.V. and Y. Lecrubier, M.I.N.I. (Mini International Neuropsychiatric Interview). Svensk översättning: Mini internationell neuropsykiatrisk intervju, Svensk reviderad version 5.0.0; Allgulander C, Waern M, Humble M, Ågren H. 2000: University of South Florida - Tampa, Hôpital de la Salpêtrière - Paris.
4. Kroenke, K., R.L. Spitzer, and J.B. Williams, The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med*, 2001. 16(9): p. 606-13.
5. Spitzer, R.L., K. Kroenke, and J.B. Williams, Validation and utility of a self-report version of PRIME-MD: the PHQ primary care study. *Primary Care Evaluation of Mental Disorders. Patient Health Questionnaire. Jama*, 1999. 282(18): p. 1737-44.
6. Svanborg, P. and M. Åsberg, A new self-rating scale for depression and anxiety states based on the Comprehensive Psychopathological Rating Scale. *Acta Psychiatr Scand*, 1994. 89(1): p. 21-8.
7. Mattila-Evenden, M., et al., Determinants of self-rating and expert rating concordance in psychiatric out-patients, using the affective subscales of the CPRS. *Acta Psychiatr Scand*, 1996. 94(6): p. 386-96.
8. Hirschfeld, R.M., et al., Development and validation of a screening instrument for bipolar spectrum disorder: the Mood Disorder Questionnaire. *Am J Psychiatry*, 2000. 157(11): p. 1873-5.
9. Young, R.C., et al., A rating scale for mania: reliability, validity and sensitivity. *Br J Psychiatry*, 1978. 133: p. 429-35.

Läkemedelsbehandling av depression

1. Paykel 1998
2. Fagiolini et al, 2001
3. Katz et al, 1997

Läkemedelsbehandling av bipolär sjukdom

1. Strakowski et al 2001

Psykologisk behandling vid depression och bipolär sjukdom

1. SBU (2004). Behandling av depressionssjukdomar: En systematisk litteraturöversikt. Volym 3. Stockholm, Statens beredning för medicinsk utvärdering.
2. Beck, A. T. (1967). Depression: Clinical, experimental, and theoretical aspects. New York, Hoeber.
3. Beck, A. T., A. J. Rush, et al. (1979). Cognitive therapy of depression. New York, Guilford Press.
4. Segal, Z. V., J. M. Williams, et al. (2002). Mindfulness-Based Cognitive Therapy for Depression. A New Approach for Preventing Relapse. New York, The Guilford Press.
5. Teasdale, J. D., Z. V. Segal, et al. (2000). "Prevention of relapse/recurrence in major depression by mindfulness based cognitive therapy." *Journal of Consulting and Clinical Psychology* 68(4): 615-23.

6. Jacobson, N. S., C. R. Martell, et al. (2001). "Behavioral activation treatment for depression: Returning to contextual roots." *Clinical Psychology: Science and Practice* 8: 255-270.
7. Dimidjian, S., K. S. Dobson, et al. (2006). "Randomized Trial of Behavioral Activation, Cognitive Therapy, and Antidepressant Medication in the Acute Treatment of Adults With Major Depression." *Journal of Consulting and Clinical Psychology* 74(4): 658-670.
8. McCullough, J. P. (2003). "Treatment for Chronic Depression Using Cognitive Behavioral Analysis System of Psychotherapy (CBASP)." *Journal of Clinical Psychology* 59: 833-846.
9. Keller, M. B., J. P. McCullough, et al. (2000). "A comparison of nefazodone, the cognitive-behavioral analysis system of psychotherapy, and their combination for the treatment of chronic depression." *New England Journal of Medicine* 343(20): 1462-70.
10. Nemeroff, C. B., C. M. Heim, et al. (2003). "Differential responses to psychotherapy versus pharmacotherapy in patients with chronic forms of major depression and childhood trauma." *PNAS* 100(24): 14293-14296.
11. McKendree-Smith, N. L., M. Floyd, et al. (2003). "Self-administered treatments for depression: a review." *Journal of Clinical Psychology* 59: 275-288.
12. Klerman, G. L., M. M. Weissman, et al. (1984). *Interpersonal Psychotherapy of Depression*. New York, Basic Books.
13. Samrådsforum (2006). *Psykoterapi. Information om olika psykoterapiinriktningar*. Stockholm, 2006.
14. Knekt, P. and O. Liljefors (2004). A randomized trial of the effect of four forms of psychotherapy on depressive and anxiety disorders: Design, methods, and results on the effectiveness of short-term psychodynamic psychotherapy and solution-focused therapy during a one-year follow-up study. Helsinki Psychotherapy Study. Helsingfors, Kela, The social insurance institution. 77: 1-115.
15. Maina, G., F. Forner, et al. (2005). "Randomized Controlled Trial Comparing brief Dynamic and Supportive Therapy with Waiting List Condition in Minor Depressive Disorders." *Psychotherapy and Psychosomatics* 74: 243-50.

16. Fava, G. A., C. Ruini, et al. (2004). "Six-Year Outcome of Cognitive Behavior Therapy for Prevention of Recurrent Depression." *American Journal of Psychiatry* 161: 1872-1876.
17. Hollon, S. D., R. J. DeRubeis, et al. (2005). "Prevention of Relapse Following Cognitive Therapy vs Medications in Moderate to Severe Depression." *Archives of General Psychiatry* 62: 417-422.
18. Scott, J. (2006). "Psychotherapy for bipolar disorders - efficacy and effectiveness." *Journal of Psychopharmacology* 20(2): 46-50.
19. Scott, J., F. Colom, et al. (2006). "A meta-analysis of relapse rates with adjunctive psychological therapies compared to usual psychiatric treatment for bipolar disorders." *International Journal of Neuropsychopharmacology*: 1-7.
20. Lam, D. H., P. Hayward, et al. (2005). "Relapse Prevention in Patients with Bipolar Disorder: Cognitive Therapy Outcome After 2 Years." *American Journal of Psychiatry* 162(2): 324-329.
21. Jones, S. (2004). "Psychotherapy of bipolar disorders: a review." *Journal of Affective Disorders* 80(2-3): 101-14.
22. Beck, J. S. and C. F. Newman (2005). *Psychotherapy for Bipolar Disorder: Treatments to Enhance Medication Adherence and Improve Outcomes.*, Medscape & Mental Health. 2005.
23. Thase, M. D. (2006). *Examining the Role of Psychotherapy in the Management of Bipolar Disorder*, Medscape Psychiatry & Mental Health. 2006.
24. Lam, D. H., S. H. Jones, et al. (2000). *Cognitive Therapy for Bipolar Disorder: A Therapist's Guide to Concepts, Methods and Practice*, John Wiley & Sons.
25. Miklowitz, D. J., T. L. Simoneau, et al. (2000). "Family-focused treatment of bipolar disorder: 1-year effects of psychoeducational programme in conjunction with pharmacotherapy." *Biological Psychiatry* 48: 582-592.
26. Weber Rouget, B. and J.-M. Aubry (2006). "Efficacy of psychoeducational approaches on bipolar disorders: A review of the literature." *Journal of Affective Disorders* In press.

Fysikaliska behandlingsmetoder

1. Ljusterapi vid depression samt övrigt behandling av årstidsbunden depression. En systematisk litteraturoversikt, SBU 2007
2. Wassermann & Lisanby, 2001
3. George et al 2000
4. Sackeim et al 2001 b

Omvårdnad

1. Ottosson, J-O. (2000) Psykiatri. 5:e uppl. Stockholm. Liber.
2. Arvidsson B. och Skärsäter I. (red.) Psykiatrisk omvårdnad – att stödja hälsobefrämjande processer. Danmark. Narayana Press.
3. Videbeck, L. (2001). Psychiatric Mental Helth Nursing s.337). Philadelphia. Lippincott Williams & Wilkins
4. Løkensgard, I. (1997) Psykiatrisk vård och specifik omvårdnad. Lund. Studentlitteratur.
5. Rawlins, R.P. (1993) Hope-Hopelessness i Rawlins, R.P., Williams, S. R., Beck, C.K. (red.) Mental health – Psychiatric nursing. A holistic life-cycle approach. 3 ed. St. Louis. Mosby-Year Book.
6. Moore, S.L. (2005). Hope makes a difference. Journal of Psychiatric and Mental Health Nursing. 12, 100-105.
7. Townsend, M. C. (1994). Nursing diagnoses in Psychiatric Nursing: a pocket guide for care plan constructions. Philadelphia. F.A. Davis company.
8. Shives, L.R. & Isaacs, A. (2002). Basic Concepts of Psychiatric-Mental Health Nursing. 5th ed. Philadelphia. Lippincott.
9. Hedman, Ahlström, B. (2006) Familjens betydelse för psykisk hälsa och ohälsa vid depression, i Arvidsson B. och Skärsäter I. (red.) Psykiatrisk omvårdnad – att stödja hälsobefrämjande processer. Danmark. Narayana Press.

10. Östman, M., Hansson, L. (2004). Appraisal of caregiving, burden and psychological distress in relatives of psychiatric inpatients. *European Psychiatry*. 19, 402-407.
11. Gutiérrez-Maldonado, J., Caqueo-Urizar, A. & Kavanagh, D.J. (2005). Burden of care and general health in families of patients with schizophrenia. *Social Psychiatry and Psychiatric Epidemiology*. 40, (11), 899-904.
12. Koukia, E. Madianos, M.G. (2005). Is psychosocial rehabilitation of schizophrenic patients preventing family burden? A comparative study. *Journal of Psychiatric and Mental Health Nursing*. 12, (4), 415-422.
13. Östman, M., Wallsten, T., Kjellin, L. (2005). Family burden and relative's participation in psychiatric care: Are the patient's diagnosis and the relation to the patient of importance? *International Journal of Social Psychiatry*. 51, 291-301.
14. Kralik, D., Kock, T., Price, K. & Howard, N (2004). Chronic illness and self-management: Talking action to create order. *Journal of clinical nursing*. Feb;13,(2), 259-267.
15. McCann, T.V. & Clark, E. (2004). Advancing self-determination with young adults who have schizophrenia. *Journal of Psychiatric and Mental Health Nursing*. 11 (1), 12-20.
16. Olofsson, B. (2000). Use of coercion in psychiatric care, as narrated by patients, nurses and physicians. Akademisk avhandling. Umeå. Umeå Universitet.
17. Hamilton, P. (Red):Psykiatrins juridik 2007. Juris AB, Stockholm 2007.
18. Svensson, J. (1996.) Patienters erfarenhet av tvångsvård. FoU- enheten/psyk, Västra Stockholms sjukvårdsområde, RSMH.
19. Haglund, K. Von Knorring, L. Von Essen, L. (2003). Forced medication in psychiatric care: Patient's experiences and nurse perceptions. *Journal of Psychiatric and Mental Health Nursing*. 10,(1):65-62.
20. RSMH:s Rättsliga råd. (2000)

Den terapeutiska alliansen

1. Bordin, 1979
2. Frank, 1991

Affektiv sjukdom under graviditet och postpartum

1. Kendell R, Chalmers J, Platz C. Epidemiology of puerperal psychoses [published erratum appears in Br J Psychiatry 1987 Jul;151:135]. Br J Psychiatry 1987;150:662-673.
2. Evans J, Heron J, Francomb H, Oke S, Golding J. Cohort study of depressed mood during pregnancy and after childbirth. Bmj 2001;323:257-60.
3. Pfulmann B, Stoeber G, Beckmann H. Postpartum psychoses: prognosis, risk factors, and treatment. Curr Psychiatry Rep 2002;4:185-90.
4. Akiskal HS, Bourgeois ML, Angst J, Post R, Moller H, Hirschfeld R. Re-evaluating the prevalence of and diagnostic composition within the broad clinical spectrum of bipolar disorders. J Affect Disord 2000;59 Suppl 1:S5-S30.
5. Rubinow DR. Antidepressant Treatment During Pregnancy: Between Scylla and Charybdis 10.1176/appi.ajp.163.6.954. American Journal of Psychiatry 2006;163:954-956.
6. Goodwin GM. Evidence-based guidelines for treating bipolar disorder: recommendations from the British Association for Psychopharmacology. J Psychopharmacol 2003;17:149-73; discussion 147.
7. Nonacs R, Cohen LS, Viguera AC, Mogielnicki J. Diagnosis and Treatment of Mood and Anxiety Disorders in Pregnancy. In: Cohen LS, Nonacs R, eds. Mood and Anxiety Disorders During Pregnancy and Postpartum. 24, number 4 vol. Washington DC: American Psychiatric Publishing.; 2005.:17-51.
8. Field T. Infants of depressed mothers. Infant Behavior and Development 1995;Volume 18:1-13.

9. Lindahl V, Pearson JL, Colpe L. Prevalence of suicidality during pregnancy and the postpartum. *Arch Women Ment Health* 2005;8:77-87.
10. Oates M. Suicide: the leading cause of maternal death. *Br J Psychiatry* 2003; 183:279-281.
11. Spinelli MG. A Systematic Investigation of 16 Cases of Neonaticide. *Am J Psychiatry* 2001;158:811-813.
12. Einarson TR, Einarson A. Newer antidepressants in pregnancy and rates of major malformations: a meta-analysis of prospective comparative studies. *Pharmacoepidemiol Drug Saf* 2005.
13. Spigset O, Hägg S. Nya antidepressiva läkemedel under graviditet och amning. *Läkartidningen* 2004;101:1176-1181.
14. Källén B. Antidepressiv behandling under graviditet - finns det risker för fostret? *SERIP* 2004:4-7.
15. Lattimore KA, Donn SM, Kaciroti N, Kemper AR, Neal CR, Jr., Vazquez DM. Selective serotonin reuptake inhibitor (SSRI) use during pregnancy and effects on the fetus and newborn: a meta-analysis. *J Perinatol* 2005;25:595-604.
16. GlaxoSmithKlein. Viktig information till förskrivare. Solna; 2005.
17. Läkemedelsverket. Seroxat (paroxetin) och risken för fosterskada. Uppsala: Läkemedelsverket; 2005:89.
18. Kallen B, Otterblad Olausson P. Antidepressant drugs during pregnancy and infant congenital heart defect. *Reprod Toxicol* 2006;21:221-2.
19. Sanz EJ, De-las-Cuevas C, Kiuru A, Bate A, Edwards R. Selective serotonin reuptake inhibitors in pregnant women and neonatal withdrawal syndrome: a database analysis. *Lancet* 2005;365:482-7.
20. Chambers CD, Hernandez-Diaz S, Van Marter LJ et al. Selective Serotonin-Reuptake Inhibitors and Risk of Persistent Pulmonary Hypertension of the Newborn 1056/NEJMoa052744. *N Engl J Med* 2006;354:579-587.
21. Adler M, von Schreeb S, Lindemalm S, Böttiger Y. SSRI-behandling under graviditet och amning - riktlinjer inom Karolinska, Huddinges upptagningsområde. Kvinnokliniken, Barnens sjukhus, Psykiatriska kliniken SV samt Klinisk Farmakaologi vid Karolinska Universitetssjukhuset i Huddinge.; 2005.
22. Nulman I, Rovet J, Stewart DE et al. Child Development Following Exposure to Tricyclic Antidepressants or Fluoxetine Throughout Fetal

- Life: A Prospective, Controlled Study 10.1176/appi.ajp.159.11.1889. American Journal of Psychiatry 2002;159:1889-1895.
23. Misri S, Reebye P, Kendrick K et al. Internalizing Behaviors in 4-Year-Old Children Exposed in Utero to Psychotropic Medications 10.1176/appi.ajp.163.6.1026. American Journal of Psychiatry 2006;163:1026-1032.
 24. Einarson A, Fatoye B, Sarkar M et al. Pregnancy outcome following gestational exposure to venlafaxine: a multicenter prospective controlled study. Am J Psychiatry 2001;158:1728-30.
 25. Ilett KF, Kristensen JH, Hackett LP, Paech M, Kohan R, Rampono J. Distribution of venlafaxine and its O-desmethyl metabolite in human milk and their effects in breastfed infants. Br J Clin Pharmacol 2002;53:17-22.
 26. McElhatton PR, Garbis HM, Elefant E et al. The outcome of pregnancy in 689 women exposed to therapeutic doses of antidepressants. A collaborative study of the European Network of Teratology Information Services (ENTIS). Reprod Toxicol 1996;10:285-94.
 27. Hendrick V, Altshuler L. Management of Major Depression During Pregnancy 1176/appi.ajp.159.10.1667. Am J Psychiatry 2002;159:1667-1673.
 28. Heikkinen T, Ekblad U, Laine K. Transplacental transfer of amitriptyline and nortriptyline in isolated perfused human placenta. Psychopharmacology (Berl) 2001;153:450-4.
 29. Wisner KL, Perel JM, Findling RL. Antidepressant treatment during breast-feeding. Am J Psychiatry 1996;153:1132-7.
 30. Källén B, Källén K. Läkemedel och fosterskador. JANUS. Läkemedelsinformation för sjukvården.; 2004.
 31. Yonkers KA, Wisner KL, Stowe Z et al. Management of Bipolar Disorder During Pregnancy and the Postpartum Period. Am J Psychiatry 2004;161:608-620.
 32. Cohen LS, Friedman JM, Jefferson JW, Johnson EM, Weiner ML. A reevaluation of risk of in utero exposure to lithium. Jama 1994;271:146-50.
 33. Viguera AC, Cohen LS, Nonacs R, Baldessarini R. Management of Bipolar Disorder During Pregnancy and the Postpartum Period: Weighing the Risks and Benefits. In: Cohen LS, Nonacs R, eds. Mood and Anxiety Disorders During Pregnancy and Postpartum. 24 vol. Washington D.C.: American Psychiatric Publishing, Inc.; 2005:53-76.

34. Viguera AC, Newport JD, Ritchie JC et al. Lithium and Lactation. Sixth International Conference on Bipolar Disorder. Pittsburgh, USA.; 2005.
35. Morrow J, Russell A, Guthrie E et al. Malformation risks of antiepileptic drugs in pregnancy: a prospective study from the UK Epilepsy and Pregnancy Register 1136/jnnp.2005.074203. *J Neurol Neurosurg Psychiatry* 2006;77:193-198.
36. Läkemedelsverket. Lamotrigin och risken för läppgomspalt. Uppsala; 2006.
37. Chaudron LH, Jefferson JW. Mood stabilizers during breastfeeding: a review. *J Clin Psychiatry* 2000;61:79-90.
38. Viguera AC, Cohen LS, Baldessarini RJ, Nonacs R. Managing bipolar disorder during pregnancy: weighing the risks and benefits. *Can J Psychiatry* 2002;47:426-36.
39. Eli-Lilly. Olanzapine - use in pregnant and nursing women, update until dec 2004.; 2005.
40. McKenna K, Koren G, Tetelbaum M et al. Pregnancy outcome of women using atypical antipsychotic drugs: a prospective comparative study. *J Clin Psychiatry* 2005;66:444-9; quiz 546.
41. Diav-Citrin O, Shechtman S, Ornoy S et al. Safety of haloperidol and penfluridol in pregnancy: a multicenter, prospective, controlled study. *J Clin Psychiatry* 2005;66:317-22.
42. Winans EA. Antipsychotics and breastfeeding. *J Hum Lact* 2001;17:344-7.
43. Croke S, Buist A, Hackett LP, Ilett KF, Norman TR, Burrows GD. Olanzapine excretion in human breast milk: estimation of infant exposure. *Int J Neuropsychopharmacol* 2002;5:243-7.
44. Cohen LS, Altshuler LL, Harlow BL et al. Relapse of major depression during pregnancy in women who maintain or discontinue antidepressant treatment. *Jama* 2006;295:499-507.
45. Viguera AC, Nonacs R, Cohen LS, Tondo L, Murray A, Baldessarini RJ. Risk of Recurrence of Bipolar Disorder in Pregnant and Nonpregnant Women After Discontinuing Lithium Maintenance. *Am J Psychiatry* 2000;157:179-184.
46. Baldessarini RJ, Tondo L, Floris G, Rudas N. Reduced morbidity after gradual discontinuation of lithium treatment for bipolar I and II disorders: a replication study. *Am J Psychiatry* 1997;154:551-3.
47. Tryckfrihetsförordning 1949:105 och sekretesslag 1980:100.; 1980.

48. Cohen LS, Sichel DA, Robertson LM, Heckscher E, Rosenbaum JF. Postpartum prophylaxis for women with bipolar disorder. *Am J Psychiatry* 1995;152:1641-5.
49. Nonacs R. Postpartum Mood Disorders. In: Cohen LS, Nonacs R, eds. *Mood and Anxiety Disorders During Pregnancy and Postpartum*. 24 vol. Washington D.C.: American Psychiatric Publishing, Inc.; 2005:77-103.
50. Wickberg B, Hwang P. [Do not minimize signs of postpartum depression! Early intervention essential to prevent negative consequences for the child]. *Lakartidningen* 2001;98:1534-8.
51. Peindl KS, Wisner KL, Hanusa BH. Identifying depression in the first postpartum year: guidelines for office-based screening and referral. *J Affect Disord* 2004;80:37-44.
52. Cox JL, Holden JM, Sagovsky R. Edinburgh Postnatal Depression Scale (EPDS); 1997.
53. Montgomery S, Åsberg M. Montgomery-Asberg Depression Rating Scale (MADRS); 1979.
54. Svanborg P, Asberg M. A comparison between the Beck Depression Inventory (BDI) and the self-rating version of the Montgomery Asberg Depression Rating Scale (MADRS). *J Affect Disord* 2001;64:203-16.
55. Spitzer RL, Kroenke K, Williams JBW, and the Patient Health Questionnaire Primary Care Study Group. Validation and Utility of a Self-report Version of PRIME-MD: The PHQ Primary Care Study. *JAMA* 1999;282:1737-1744.
56. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001;16:606-13.
57. Pertrillo LF, Nonacs R, Viguera AC, Cohen LS. Course of Psychiatric Illness During Pregnancy and the Postpartum. In: Cohen LS, Nonacs R, eds. *Mood and Anxiety Disorders During Pregnancy and Postpartum*. 24 vol. Washington D.C.: American Psychiatric Publishing, Inc.; 2005:1-15.

Bipolär sjukdom hos barn och unga

1. Backlund Lena (2004) Tidig upptäckt av bipolär sjukdom – en viktig angelägenhet för hela sjukvården, Incitament Nr 1:507-11
2. Geller Barbara, Del Bello, Melissa P (2003) Bipolär disorders in childhood and early dolescence, Guilford Press, New York
3. Olsson Gunilla (2004) Bipolär sjukdom hos barn och tonåringar- en diagnos under debatt, Läkartidningen, Vol:101 Nr 13; 1182-1187
4. Pavaluri Mani N, Birmaher Boris, Naylor Michael W, Pediatric bipolar disorder; A review of the past 10 years. J Am Acad Child Adolesc Psychiatry, 44:9, sept 2005, 846-871

Affektiv sjukdom hos personer över 65 år

1. Gottfries, 1998
2. Gottfries & Karlsson, 2000

Beställning

Kontorsservice

Tel: 08-737 49 57; Fax: 08-737 49 59

E-post: informationsmaterial.lsf@sll.se

Postadress: Box 6401, 113 82 Stockholm

Besöksadress: Olivecronas väg 7, bv; Sabbatsbergs sjukhus

Rapporter från Medicinskt programarbete är publicerade på:
www.webbhotell.sll.se/sv/cvu/Medicinska-program-och-kunskapsstod



Stockholms läns landsting

Forum för kunskap och gemensam utveckling

Box 12175, 118 91 Stockholm

Telefon 08-737 30 00. Fax 08-737 44 64